arXiv:1106.0422v2 [math.GT] 3 Jun 2011

On upper bounds on stable commutator lengths in mapping class groups

Naoyuki Monden

Abstract

We give new upper bounds on the stable commutator lengths of Dehn twists in mapping class groups and new lower bounds on the stable commutator lengths of Dehn twists in hyperelliptic mapping class groups. In particular, we show that the stable commutator lengths of Dehn twists about a nonseparating and a separating curve on an oriented closed surface of genus 2 are not equal to each other.

1 Introduction

Let G be a group, and let $[G,G]$ denote the commutator subgroup of G. Given $x \in [G,G]$ the *commutator length* $\mathrm{cl}_G(x)$ of x is the least number of commutators in G whose product is equal to x. The *stable commutator length* $\mathrm{scl}_G(x)$ is the limit of $\mathrm{cl}_G(x^n)/n$ as n goes to infinity. If $x^m \in [G,G]$ for some positive integer m, define $\mathrm{scl}_G(x) = \mathrm{scl}_G(x^m)/m$, and define $\mathrm{scl}_G(x) = \infty$ if no power of x is contained in $[G,G]$ (We refer the reader to [Ca] for the details of the theory of the stable commutator length).

Let Σ_g be a closed connected oriented surface of genus $g \geq 2$ embedded in $\mathbb{R}^3$ in Figure 1. We can define the hyperelliptic involution $\iota : \Sigma_g \to \Sigma_g$ as in Figure 1.



Figure 1: hyperelliptic involution ι and the curves $s_1, \dots s_{g-1}$

Let $\mathcal{M}_g$ be the mapping class group of Σ_g, i.e., the group of isotopy classes of orientation-preserving diffeomorphisms of Σ_g. Let t_c and t_s denote the right handed Dehn twist about a nonseparating curve c and a nontrivial separating curve s on Σ_g, respectively.

It is well known that $\mathcal{M}_g/[\mathcal{M}_g, \mathcal{M}_g]$ is generated by the class of a Dehn twist about a nonseparating simple closed curve and is equal to $\mathbb{Z}_{10}$ if $g = 2$ and trivial if $g \geq 3$. Therefore, If $g \geq 3$, then t_c and t_s are in $[\mathcal{M}_g, \mathcal{M}_g]$. Moreover, it is well known that t_c^{10} and t_s^5 are in $[\mathcal{M}_2, \mathcal{M}_2]$. Hence, for any $g \geq 2$ we can define $\mathrm{scl}_{\mathcal{M}_g}(t_c)$ and $\mathrm{scl}_{\mathcal{M}_g}(t_s)$.

Endo and Kotschick proved that $\mathrm{scl}_{\mathcal{M}_g}(t_s) \geq 1/(18g-6)$ (see [EK]). Consequently, they proved that for any $g \geq 2$ $\mathcal{M}_g$ is not uniformly perfect and that the natural map from the second bounded cohomology to the ordinary cohomology of $\mathcal{M}_g$ is not injective, which verified two conjectures of Morita [Mo]. Korkmaz proved that $\mathrm{scl}_{\mathcal{M}_g}(t_c) \geq 1/(18g-6)$ (see [Ko]). He also gave upper bounds for the stable commutator lengths of Dehn twists. He showed that (1) $\mathrm{scl}_{\mathcal{M}_g}(t_c) \leq 3/20$ for $g \geq 2$ and (2) $\mathrm{scl}_{\mathcal{M}_g}(t_s) \leq 3/4$ for $g \geq 3$ by using results of [Ba] and [Cu] and by showing that t_c^{10} is written as a product of two commutators. By using the results of quasimorphisms we give the following upper bounds.

Theorem 1.1. *Let c and s be a nonseparating curve and a separating curve on Σ_g $(g \geq 2)$, respectively.*

(1) *If $g \geq 2$, then* $\mathrm{scl}_{\mathcal{M}_g}(t_c) \leq \dfrac{1}{10}$,

(2) *If $g \geq 3$, then* $\mathrm{scl}_{\mathcal{M}_g}(t_s) \leq \dfrac{1}{2}$,

(3) *If $g = 2$, then* $\mathrm{scl}_{\mathcal{M}_2}(t_s) \leq \dfrac{7}{10}$.

Let $\mathcal{H}_g$ be the hyperelliptic mapping class group of genus g, i.e., the subgroup of $\mathcal{M}_g$ which consists of all isotopy classes of orientation-preserving diffeomorphisms of Σ_g commuting with the isotopy class of ι. Let $s_1, \ldots, s_{g-1}$ be separating curves as shown in Figure 1. Each t_{s_h} $(h = 1, \ldots, g)$ belongs to $\mathcal{H}_g$ and $t_{s_{g-h}}$ is conjugate to t_{s_h} in $\mathcal{H}_g$. Since the stable commutator length is constant on conjugacy classes, it suffices to consider t_{s_h} $(h = 1, \ldots, [\frac{g}{2}])$. It is well known that $t_c^{4(2g+1)}$ and $t_{s_h}^{4(2g+1)}$ are in $[\mathcal{H}_g, \mathcal{H}_g]$. Hence, we can define $\mathrm{scl}_{\mathcal{H}_g}(t_c)$ and $\mathrm{scl}_{\mathcal{H}_g}(t_{s_h})$.

Endo and Kotschick proved that $\mathrm{scl}_{\mathcal{H}_g}(t_{s_h}) \geq 1/(18g-6)$ (see [EK]). We give the following lower bounds on the stable commutator lengths of Dehn twists in $\mathcal{H}_g$.

Theorem 1.2. *For all $g \geq 2$,*

$$(1) \quad \frac{1}{4(2g+1)} \leq \mathrm{scl}_{\mathcal{H}_g}(t_c), \quad (2) \quad \frac{h(g-h)}{g(2g+1)} \leq \mathrm{scl}_{\mathcal{H}_g}(t_{s_h}) \quad (h = 1, \ldots, [\tfrac{g}{2}]).$$

In particular, Since $\mathcal{M}_2$ is equal to $\mathcal{H}_2$, by combining Theorem 1.1 with Theorem 1.2 we have $\frac{1}{20} \leq \mathrm{scl}_{\mathcal{M}_2}(t_c) \leq \frac{1}{10} \leq \mathrm{scl}_{\mathcal{M}_2}(t_s) \leq \frac{7}{10}$. On the other hand, we give the following lemma.

Lemma 1.3. $6\mathrm{scl}_{\mathcal{M}_2}(t_c) \leq \dfrac{1}{2}\mathrm{scl}_{\mathcal{M}_2}(t_s) + \dfrac{1}{2}$.

If $\mathrm{scl}_{\mathcal{M}_2}(t_c) = \mathrm{scl}_{\mathcal{M}_2}(t_s)$, then by Lemma 1.3 we have $\mathrm{scl}_{\mathcal{M}_2}(t_s) \leq \frac{1}{11}$. This contradicts to $\frac{1}{10} \leq \mathrm{scl}_{\mathcal{M}_2}(t_s)$. Therefore, we have the following results.

Corollary 1.4. *If $g = 2$, then*

(1) $\dfrac{1}{20} \leq \mathrm{scl}_{\mathcal{M}_2}(t_c) \leq \dfrac{1}{10} \leq \mathrm{scl}_{\mathcal{M}_2}(t_s) \leq \dfrac{7}{10}$.

(2) $\mathrm{scl}_{\mathcal{M}_2}(t_c)$ *is not equal to* $\mathrm{scl}_{\mathcal{M}_2}(t_s)$.

2 Preliminaries

2.1 Stable commutator lengths and quasimorphisms

Let G denote a group and let $[G,G]$ denote the commutator subgroup, the subgroup of G generated by all commutators $[x,y]=xyx^{-1}y^{-1}$ for $x,y\in G$.

Definition 2.1. *For $x\in[G,G]$, the commutator length $\mathrm{cl}_G(x)$ of x is the least number of commutators in G whose product is equal to x.*

Definition 2.2. *For $x\in[G,G]$,*

$$\mathrm{scl}_G(x)=\lim_{n\to\infty}\frac{\mathrm{cl}_G(x^n)}{n}$$

is called the stable commutator length of x.

For each fixed x, the function $n\to\mathrm{cl}_G(x^n)$ is non-negative and $\mathrm{cl}_G(x^{m+n})\leq\mathrm{cl}_G(x^m)+\mathrm{cl}_G(x^n)$; hence this limit exists. If x is not in $[G,G]$ but has a power x^m which is, define $\mathrm{scl}_G(x)=\mathrm{scl}_G(x^m)/m$. We define $\mathrm{scl}_G(x)=\infty$ if no power of x is contained in $[G,G]$.

Definition 2.3. *A quasimorphism is a function*

$$\phi: G\to\mathbb{R}$$

for which there is a least constant $D(\phi)\geq 0$ such that

$$|\phi(xy)-\phi(x)-\phi(y)|\leq D(\phi)$$

for all $x,y\in G$. We call $D(\phi)$ the defect of ϕ.

Definition 2.4. *A quasimorphism is homogeneous if it satisfies the additional property*

$$\phi(x^n)=n\phi(x)$$

for all $x\in G$ and $n\in\mathbb{Z}$.

We recall the following basic facts.

Lemma 2.5. *Let ϕ be a homogeneous quasimorphism. For all $x,y\in G$,*

(a) *$\phi(x)=\phi(yxy^{-1})$,*

(b) *$xy=yx\Rightarrow\phi(xy)=\phi(x)+\phi(y)$.*

Proof. For any positive integer n,

$$|\phi(yxy^{-1})-\phi(x)|=\frac{1}{n}|\phi(yx^ny^{-1})-\phi(x^n)|\leq\frac{2D(\phi)}{n}.$$

Hence, we have $\phi(x)=\phi(yxy^{-1})$.

Suppose that $xy=yx$. For any positive integer n,

$$\begin{aligned}|\phi(xy)-\phi(x)-\phi(y)| &= \frac{1}{n}|\phi((xy)^n)-\phi(x^n)-\phi(y^n)|\\ &= \frac{1}{n}|\phi(x^ny^n)-\phi(x^n)-\phi(y^n)|\leq\frac{1}{n}D(\phi).\end{aligned}$$

Hence, $\phi(xy)=\phi(x)+\phi(y)$.

□

Theorem 2.6 (Bavard's Duality Theorem [Ba])**.** *Let Q be the set of homogeneous quasimorphisms on G. For any $x \in [G,G]$, we have*

$$\mathrm{scl}_G(x) = \sup_{\phi \in Q} \frac{|\phi(x)|}{2D(\phi)}.$$

2.2 Relations in mapping class groups

Hereafter, we do not distinguish a simple closed curve and its isotopy class. The next lemmas are well known.

Lemma 2.7. *For any $f \in \mathcal{M}_g$ and any simple closed curve c in Σ_g we have*

$$t_{f(c)} = f t_c f^{-1}.$$

Lemma 2.8. *Let c and d be two simple closed curves on Σ_g.*
(a) *If c is disjoint from d, then $t_c t_d = t_d t_c$.*
(b) *If c intersects d in one point transversely, then $t_c t_d t_c = t_d t_c t_d$.*

The following two relations in $\mathcal{M}_g$ are also well known. The first one is the *lantern relation*. This relation was discovered by Dehn (see [De]) and was rediscovered by Johnson (see [Jo]). Let s, a, b, c, x, y and z be simple closed curves on Σ_g ($g \geq 3$) in Figure 2. The Dehn twists about these simple closed curves satisfy the *lantern relation*

$$t_s t_a t_b t_c = t_x t_y t_z.$$



Figure 2: The curves s, a, b, c, x, y and z on Σ_g ($g \geq 3$).

In the case of $g = 2$, we define the curves s, a_1, a_3, a_5 and x on Σ_2 in Figure 3. t_s, t_{a_1}, t_{a_3}, t_{a_5} and t_x satisfy the *lantern relation*

$$t_{a_1}^2 t_{a_5}^2 = t_{a_3} t_s t_x.$$



Figure 3: The curves s, a, b, c, x, y and z on Σ_2.



Figure 4: The curves $a_1 a_2$, a_3, a_4 and a_5.

The second relation is the *chain relation*. Let a_1, a_2, a_3, a_4 and a_5 be simple closed curves on Σ_g in Figure 4. t_{a_1}, t_{a_2}, t_{a_3}, t_{a_4} and t_{a_5} satisfy the *chain relation*

$$(t_{a_3} t_{a_2} t_{a_1})^4 = t_{a_4} t_{a_5}.$$

We note that if $g = 2$, then a_4 is equal to a_5. Therefore, in the case of $g = 2$, the chain relation is as follows:

$$(t_{a_3} t_{a_2} t_{a_1})^4 = t_{a_5}^2.$$

3 Proofs of the main results

We prove Theorem 1.1.

Proof of Theorem 1.1. Let c and s be a nonseparating curve and a separating curve on Σ_g, respectively. Let φ be a homogeneous quasimorphism on $\mathcal{M}_g$.

We first prove that $\mathrm{scl}_{\mathcal{M}_g}(t_c) \leq \frac{1}{10}$ for $g \geq 2$. Suppose that $g \geq 3$. By the chain relation and Lemma 2.8

$$\begin{array}{rcl}
t_{a_4} t_{a_5} & = & t_{a_3} t_{a_2} \underline{t_{a_1} t_{a_3}} t_{a_2} t_{a_1} t_{a_3} t_{a_2} \underline{t_{a_1} t_{a_3}} t_{a_2} t_{a_1} \\
& = & t_{a_3} t_{a_2} t_{a_3} \underline{t_{a_1} t_{a_2} t_{a_1} t_{a_3} t_{a_2} t_{a_3}} t_{a_1} t_{a_2} t_{a_1} \\
& = & t_{a_3} \underline{t_{a_2} t_{a_3} t_{a_2}} t_{a_1} t_{a_2} t_{a_2} t_{a_3} \underline{t_{a_2} t_{a_1} t_{a_2}} t_{a_1} \\
& = & t_{a_3} t_{a_3} t_{a_2} t_{a_3} t_{a_1} t_{a_2} t_{a_2} t_{a_3} t_{a_1} t_{a_2} t_{a_1} t_{a_1} \\
& = & t_{a_3}^2 (t_{a_2} t_{a_3} t_{a_1} t_{a_2})(t_{a_2} t_{a_3} t_{a_1} t_{a_2}) t_{a_1}^2.
\end{array}$$

Therefore, we have $t_{a_4} t_{a_5} t_{a_3}^{-2} t_{a_1}^{-2} = (t_{a_2} t_{a_3} t_{a_1} t_{a_2})^2$. Since a_1, a_3, a_4 and a_5 are disjoint from each other, from the definition of the homogeneous quasimorphism, Lemma 2.5, Lemma 2.7 and Lemma 2.8 we have

$$\begin{array}{rcl}
\varphi(t_{a_2} t_{a_3} t_{a_1} t_{a_2}) & = & \dfrac{1}{2}(\varphi(t_{a_4} t_{a_5} t_{a_3}^{-2} t_{a_1}^{-2})) \\
& = & \dfrac{1}{2}(\varphi(t_{a_4}) + \varphi(t_{a_5}) - 2\varphi(t_{a_3}) - 2\varphi(t_{a_1})) \\
& = & -\varphi(t_c).
\end{array}$$

On the other hand, by Lemma 2.5

$$-\varphi(t_c) = \varphi(t_{a_2} t_{a_3} t_{a_1} t_{a_2}) = \varphi(t_{a_2}(t_{a_2} t_{a_3} t_{a_1} t_{a_2}) t_{a_2}^{-1}) = \varphi(t_{a_2}^2 t_{a_3} t_{a_1}).$$

Therefore, from the definition of the quasimorphim and Lemma 2.5

$$\begin{array}{rcl}
D(\varphi) & \geq & |\varphi(t_{a_2}^2 t_{a_3} t_{a_1}) - \varphi(t_{a_2}^2) - \varphi(t_{a_3} t_{a_1})| \\
& = & |-\varphi(t_c) - 2\varphi(t_{a_2}) - \varphi(t_{a_3}) - \varphi(t_{a_1})| \\
& = & 5|\varphi(t_c)|.
\end{array}$$

Since $\frac{|\varphi(t_c)|}{2D(\varphi)} \le \frac{1}{10}$, by Bavard's Duality Theorem we have $\mathrm{scl}_{\mathcal{M}_g}(t_c) \le \frac{1}{10}$.

If $g = 2$, we use the chain relation $(t_{a_3}t_{a_2}t_{a_1})^4 = t_{a_5}^2$. By a similar argument we can prove that $\mathrm{scl}_{\mathcal{M}_2}(t_c) \le \frac{1}{10}$.

We next prove that $\mathrm{scl}_{\mathcal{M}_g}(t_s) \le \frac{1}{2}$ for $g \ge 3$. By the lantern relation and Lemma 2.8

$$t_s t_a t_b t_c t_z^{-1} = t_x t_y.$$

Hence, by Lemma 2.8 and Lemma 2.5 we have

$$\begin{aligned} \varphi(t_x t_y) &= \varphi(t_s t_a t_b t_c t_z^{-1}) \\ &= \varphi(t_s) + \varphi(t_a) + \varphi(t_b) + \varphi(t_c) - \varphi(t_z) \\ &= \varphi(t_s) + 2\varphi(t_c). \end{aligned}$$

From the definition of quasimorphism and Lemma 2.5

$$\begin{aligned} D(\varphi) &\ge |\varphi(t_x t_y) - \varphi(t_x) - \varphi(t_y)| \\ &= |\varphi(t_s) + 2\varphi(t_c) - \varphi(t_x) - \varphi(t_y)| \\ &= |\varphi(t_s)|. \end{aligned}$$

By Bavard's Duality Theorem we have $\mathrm{scl}_{\mathcal{M}_g}(t_s) \le \frac{1}{2}$.

Finally, we prove that $\mathrm{scl}_{\mathcal{M}_2}(t_s) \le 7/10$. By the lantern relation we have $t_{a_1}^2 t_{a_5}^2 t_{a_3}^{-1} = t_s t_x$. From the definition of quasimorphism and Lemma 2.5

$$\begin{aligned} D(\varphi) &\ge |\varphi(t_s t_x) - \varphi(t_s) - \varphi(t_x)| \\ &= |\varphi(t_{a_1}^2 t_{a_5}^2 t_{a_3}^{-1}) - \varphi(t_s) - \varphi(t_x)| \\ &= |2\varphi(t_{a_1}) + 2\varphi(t_{a_5}) - \varphi(t_{a_3}) - \varphi(t_s) - \varphi(t_x)| \\ &= |2\varphi(t_c) - \varphi(t_s)| \\ &\ge |\varphi(t_s)| - 2|\varphi(t_c)|. \end{aligned}$$

Therefore, we have $\frac{|\varphi(t_s)|}{2D(\varphi)} \le 2\frac{|\varphi(t_c)|}{2D(\varphi)} + \frac{1}{2}$. When we use Bavard's duality theorem for the left side after having used Bavard's duality theorem for the right side, we have $\mathrm{scl}_{\mathcal{M}_2}(t_s) \le 2\mathrm{scl}_{\mathcal{M}_2}(t_c) + \frac{1}{2}$. Since $\mathrm{scl}_{\mathcal{M}_2}(t_c) \le \frac{1}{10}$, $\mathrm{scl}_{\mathcal{M}_2}(t_s) \le \frac{7}{10}$.

This completes the proof of Theorem 1.1. □

Let Σ_k be a closed connected oriented surface of genus $k \ge 1$. We denote the signature of a 4-manifold M as $\sigma(M)$ in the rest of this paper.

Theorem 3.1 ([Ma1],[Ma2],[E]). *Let M be a 4-manifold which admits a hyperelliptic Lefschetz fibration of genus g over Σ_k. Let n and $s = \Sigma_{h=1}^{[\frac{g}{2}]} b_h$ be the numbers of nonseparating and separating vanishing cycles in the global monodromy of this fibration, respectively. Then*

$$\sigma(M) = -\frac{g+1}{2g+1}n + \Sigma_{h=1}^{[\frac{g}{2}]}(\frac{4h(g-h)}{2g+1} - 1)b_h,$$

where b_h denotes the number of separating vanishing cycles which separate the genus g surface into two surfaces one of which has genus h.

We prove Theorem 1.2.

Proof of Theorem 1.2. We base on the argument of [Ko]. Suppose that $g \geq 3$. We assume the contrary that $\mathrm{scl}_{\mathcal{H}_g}(t_c) < \frac{1}{4(2g+1)}$. Choose a rational number r with $\mathrm{scl}_{\mathcal{H}_g}(t_c) < r < \frac{1}{4(2g+1)}$. Then there exists an arbitrarily large positive integer n such that rn is an integer and t_c^n can be written as a product of rn commutators in $\mathcal{H}_g$. Note that we take n as a multiple of $4(2g+1)$. This gives a relatively minimal genus-g Lefschetz fibration $M \to \Sigma_{rn}$ with the nonseparating vanishing cycle c repeated n times. (More details of the theory of Lefschetz fibrations can be found in [GS]).

In [Ko], Korkmaz gave an upper bound for the signature of M;

$$\sigma(M) \leq 4grn - n + 4.$$

On the other hand, by Theorem 3.1 we see

$$\sigma(M) = -\frac{g+1}{2g+1}n.$$

Hence, we obtain

$$-\frac{g+1}{2g+1}n \leq 4grn - n + 4.$$

As a result of this, we conclude that there exists arbitrarily big n such that

$$0 \leq (4r - \frac{1}{2g+1})gn + 4.$$

Since $r - \frac{1}{4(2g+1)}$ is negative, this contradicts to the inequality. This completes the proof of $\frac{1}{4(2g+1)} \leq \mathrm{scl}_{\mathcal{H}_g}(t_c)$. By a similar argument we can prove that $\frac{h(g-h)}{g(2g+1)} \leq \mathrm{scl}_{\mathcal{H}_g}(t_{s_h})$ $(h = 1, \ldots, \frac{[g]}{2})$ for $g \geq 3$. This completes the proof of Theorem 1.2. □

Finally, we will show Lemma 1.3.

Proof of Lemma 1.3. Let s, a_1 and a_2 be simple closed curves in Figure 3. It is well known that t_s, t_{a_1} and t_{a_2} satisfy the following relation.

$$t_s = (t_{a_2}t_{a_1})^6.$$

By Lemma 2.8

$$\begin{aligned}
t_s &= \underline{t_{a_2}t_{a_1}t_{a_2}t_{a_1}t_{a_2}t_{a_1}}t_{a_2}t_{a_1}\underline{t_{a_2}t_{a_1}t_{a_2}}t_{a_1} \\
&= t_{a_1}\underline{t_{a_2}t_{a_1}t_{a_2}}t_{a_1}t_{a_2}t_{a_2}t_{a_1}t_{a_1}t_{a_2}t_{a_1}t_{a_1} \\
&= t_{a_1}t_{a_1}t_{a_2}t_{a_1}t_{a_1}t_{a_2}t_{a_2}t_{a_1}t_{a_1}t_{a_2}t_{a_1}t_{a_1}
\end{aligned}$$

Therefore, from Lemma 2.8 we have $t_s t_{a_1}^{-4} = (t_{a_2}t_{a_1}t_{a_1}t_{a_2})^2$.

Let φ be a homogeneous quasimorphism on $\mathcal{M}_2$. By the definition of homogeneous quasimorphisms and Lemma 2.5 we have

$$\frac{1}{2}\varphi(t_s) - 2\varphi(t_{a_1}) = \varphi(t_{a_2}t_{a_1}t_{a_1}t_{a_2}) = \varphi(t_{a_2}(t_{a_2}t_{a_1}t_{a_1}t_{a_2})t_{a_2}^{-1}) = \varphi(t_{a_2}^2 t_{a_1}^2).$$

Let c be a nonseparating curve on Σ_2. From the definition of quasimorphisms and homogeneous quasimorphisms

$$\begin{aligned} D(\varphi) &\geq |\varphi(t_{a_2}^2 t_{a_1}^2) - \varphi(t_{a_2}^2) - \varphi(t_{a_1}^2)| \\ &= |\varphi(t_{a_2}^2 t_{a_1}^2) - 2\varphi(t_{a_2}) - 2\varphi(t_{a_1})| \\ &= |\frac{1}{2}\varphi(t_s) - 2\varphi(t_{a_1}) - 2\varphi(t_{a_2}) - 2\varphi(t_{a_1})| \\ &= |\frac{1}{2}\varphi(t_s) - 6\varphi(t_c)| \\ &\geq 6|\varphi(t_c)| - \frac{1}{2}|\varphi(t_s)|. \end{aligned}$$

Therefore, we have $6\frac{|\varphi(t_c)|}{2D(\varphi)} \leq \frac{1}{2}\frac{|\varphi(t_s)|}{2D(\varphi)} + \frac{1}{2}$. When we use Bavard's duality theorem for the left side after having used Bavard's duality theorem for the right side, we have

$$6\mathrm{scl}_{\mathcal{M}_2}(t_c) \leq \frac{1}{2}\mathrm{scl}_{\mathcal{M}_2}(t_s) + \frac{1}{2}.$$

This completes the proof of Lemma 1.3. ☐

Acknowledgment

The author would like to thank Andrew Putman for pointing out the mistake of references in this paper.

Reference

[Ba] C. Bavard, *Longueur stable des commutateurs*, L'Enseignement Mathématique 37 (1991), 109–150.

[Ca] D. Calegari, *scl*, MSJ Memoirs, 20. Mathematical Society of Japan, Tokyo, 2009

[Cu] M. Culler, *Using surfaces to solve equations in free groups*, Topology 20 (1981), no. 2, 133–145

[De] M. Dehn, *Papers on group theory and topology.* Springer-Verlag, New York, 1987 (Die Gruppe der Abbildungsklassen, Acta Math. Vol. **69** (1938), 135–206)

[E] H. Endo, Meyerfs signature cocyle and hyperelliptic fibrations, Math. Ann., 316 (2000), 237–257.

[EK] H. Endo and D. Kotschick, *Bounded cohomology and non-uniform perfection of mapping class groups*, Invent. Math. 144 (2001), no. 1, 169–175.

[GS] R. E. Gompf and A. I. Stipsicz, *4-manifolds and Kirby calculus*, AMS Graduate Studies in Mathematics 20 1999.

[Jo] D. Johnson, *Homeomorphisms of a surface which act trivially on homology.* Proc. Amer. Math. Soc. 75 (1979), no. 1, 119–125.

[Ko] M. Korkmaz, *Stable commutator length of a Dehn twist*, Michigan Math. J. 52 (2004), no. 1, 23–31.

[Ma1] Y. Matsumoto, *On 4-manifolds fibered by tori II*, Proc. Japan Acad., 59A (1983), 100–103.

[Ma2] Y. Matsumoto, *Lefschetz fibrations of genus two - a topological approach*, Proceedings of the 37th Taniguchi Symposium on Topology and Teichmüller Spaces, ed. Sadayoshi Kojima et al., World Scientific (1996), 123–148. CMP 99:06.

[Mo] S. Morita, *Structure of the mapping class groups of surfaces: a survey and a prospect*, Geometry and Topology Monographs Vol. 2, Proc. of Kirbyfest 1999, 488–505.

Department of Mathematics, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan
E-mail address: **n-monden@cr.math.sci.osaka-u.ac.jp**